

Cardiodiagnostics Services Requisition

CARDIODIAGNOSTICS DEPARTMENT Phone: 519-749-6938 Fax: 519-749-6871

| PATIENT INFORMATION: | R | EFERRING PHYSICIAN: | | | |
|-------------------------------|-------------------------------------|-----------------------------------|---|--|--|
| Last Name: | First Name: | lame: | | | |
| DOB: (dd/mm/yyyy) Health | Α | address: | | | |
| Card Number: | S | treet | | | |
| Address: | C | îty Provii | nce Postal Code | | |
| Street | P | hone: | Fax: | | |
| City | Province A | additional copies: | | | |
| Postal Code | Phone: | | | | |
| Height: Weight | | | been seen by a Cardiologist: Dr | | |
| TO BOOK A TEST CALL C | ENTRAL BOOKINGS: 519-749-69 | 990 FAX NON URGEN | IT REQUISITIONS TO: 519-749-6989 | | |
| For URGENT (da | ys) requests please contact the Ca | rdiodiagnostics Departme | ent directly at 519-749-6938 | | |
| Patient Location: Home | □Hospital: | | | | |
| | rst Available Cardiologist or | | | | |
| Urgency: ☐ Days ☐ Weeks | | | | | |
| | | gery (if known): | | | |
| | Yes If yes, Specify Language | | | | |
| · | ☐ Transthroacic Echocardiogran | | | | |
| ECHOCARDIOGRAPHY | ☐ Transesophageal Echocardiog | • | (bubble study) — Contrast | | |
| Echocanbiodnai III | Transesophagear Echocardiog | iaiii | | | |
| | ☐ 12 Lead ECG | | | | |
| ELECTROCARDIOGRAPHY | Holter Monitor: ☐ 24 hour ☐ 48 hour | | | | |
| | Loop Recorder: ☐ 14 day ☐ 28 Day | | | | |
| | | | | | |
| | ☐ Stress Test only ☐ Str | ess Test with Consult | ☐ Stress Test +/- Consult | | |
| | Select appropriate test: | | | | |
| STRESS TESTING | ☐ Treadmill Stress Test (Patient | has <u>no</u> physical, cognitive | or other impediment to exercise) | | |
| | Exercise Stress Echocardiogra | m (abnormal ECG, LVH, pr | ior CABG, Digoxin use, Female*) | | |
| | *Consider stress imaging in fema | le patients, due to high fai | lse positive rate with treadmill alone | | |
| | Dobutamine Stress Echocardi | ogram | | | |
| | ☐ Ischemia (unable to exerc | cise) 🗖 Viability 🗖 Lo | ow-flow Low-gradient aortic stenosis | | |
| | ☐ Nuclear Myocardial Perfusion | Scan | | | |
| | ☐ Exercise ☐ Persantine | Rest only (for viabil | ity) 🗖 MUGA (Wall Motion/EF) | | |
| INDICATION: Check all that ap | ply ** Requisitions without appr | | | | |
| | | | sue Model: | | |
| ☐ Chest pain ☐ Dyspnea ☐ | Palpitations | pe 🗖 Murmur: | | | |
| | | | ardial Disease | | |
| | | | | | |
| ☐ Cardiac screening for asymp | tomatic patients with multiple car | rdiovascular risk factors (s | elect all that apply): | | |
| | Dyslipidemia Hypertension | | | | |
| CLINICAL INFORMATION: | /- p //- | | _ , , , , , , , , , , , , , , , , , , , | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| Physician's Signature: | | Dat | te: | | |
| Office Use Only | | | | | |
| Office Use Offig | | | | | |

Suggested Chest Pain Assessment Algorithm (Excluding Acute Coronary Syndromes)


STEP 1. Estimate Pretest Probability of Obstructive Coronary Disease as the cause for the patient's chest pain:

Chest Pain Characteristics:

- Substernal chest discomfort, with characteristic quality and duration
- 2. Provoked by exertion or emotional stress
- **3.** Relieved by rest and/or Nitroglycerine

| | Non-Anginal Chest Pain ≤ 1 of 3 | | Atypical Chest Pain 2 of 3 | | Typical Anginal Chest Pain 3 of 3 | |
|-------|---------------------------------------|--------|-------------------------------|--------|---|--------|
| Age | Male | Female | Male | Female | Male | Female |
| 30-39 | 4% | 2% | 34% | 12% | 76% | 26% |
| 40-49 | 13% | 3% | 51% | 22% | 87% | 55% |
| 50-59 | 20% | 7% | 65% | 33% | 93% | 73% |
| 60-69 | 27% | 14% | 72% | 51% | 94% | 86% |

STEP 2. Determine the appropriate non-invasive risk stratification method:



Adapted from ACC 2012 Guideline for the diagnosis and Management of Patients with Stable Ischemic Heart Disease. Circulation. 2012. 126:e354-e471.

Stress Test with Consultation & +/- Consultation Services:

- Appropriate for the evaluation of patients presenting with chest pain or dyspnea with intermediate to high pre-test probability of obstructive CAD
- Cardiovascular screening for asymptomatic patients with multiple cardiovascular risk factors
- Pre-operative cardiac assessment, in patients with multiple cardiovascular risk factors or known CAD, not currently followed by a Cardiologist, <u>WHEN</u> it will change management
- +/- Consultation means a consultation will be provided in the event of a high risk study
- Stress test with Consultation service is <u>NOT</u> appropriate for patients who are currently being followed and managed by
 a Cardiologist. In this case, either refer directly to that physician's office or order a test only, with the results copied to
 the patient's usual Cardiologist